

Management of Infective Conjunctivitis in primary care

Summary of key learning points

- Infective conjunctivitis is a common self-limiting condition
- Diagnosis is clinical
- Swabs are unnecessary in routine practice
- Topical antibiotics should not be routinely prescribed unless special circumstances require shorter symptom duration
- Education of patient, family and school is important to reduce expectation of antibiotic drops
- If symptoms persist after 10 days, stop all drops and seek an Ophthalmology opinion
- **RED FLAGS:** red sticky eyes in newborn; soft contact lens use with corneal symptoms (photophobia, watering).

Evidence base

- Little evidence that symptoms or signs distinguish between viral or bacterial causes¹
- In proven bacterial conjunctivitis meta-analysis shows faster time to cure with topical antibiotics. However 60% of signs and 40% swabs are negative in 5 days even in placebo group²
- Although self-limiting, clinically diagnosed infective conjunctivitis treated with antibiotic drops has lower symptom severity and quicker resolution^{3,4}
- Be aware that this effect is less in children⁵
- Cochrane Database Systematic Review 2012 showed no reports of serious outcomes in either the active or placebo arms of 11 reviewed trials (3673 participants)⁶
- Therefore, patients do not appear to come to harm if topical antibiotics are withheld⁶

Why is this important to GPs?

- Acute red eye is common estimated at 2-5% of all GP consultations⁷
- Acute infective conjunctivitis probably accounts for a 1/3 of all acute red eyes presenting to GPs⁸
- With over the counter Chloramphenicol available since 2005, the diagnosis may also include elements of drop allergy and preservative-related red eye
- GP should encourage supportive measures other than topical antibiotics
- Of course there are other causes of acute red eye which are managed differently

Pathophysiology and clinical stereotypes

- Infective conjunctivitis is self limiting and is thought to be caused by viruses or bacteria in almost equal frequency.
- In studies, between 30- 50% conjunctival swabs are positive for bacteria¹⁻³. The commonest reported pathogens are streptococcus pneumoniae, staphylococcus aureus and haemophilus influenzae. Because swab results do not influence disease course in most cases, bacterial culture is used only in severe or atypical disease.
- The following rules of thumbs are useful: Bacterial - glued lids on waking; Viral – usually a history of previous episodes; Allergic – itching is common¹.
- Conjunctivitis caused by sexually transmitted infections (STI) are often more severe (Gonorrhoea) or with prolonged mucopurulent discharge (Chlamydia) and require swabs, contact tracing and treatment.
- **Red Flag**. Neonatal (less than 4 weeks) sticky eye **with redness** must be referred for specialist care urgently. The neonatal cornea is vulnerable to penetration by potential STI organisms. Sticky eye without redness is managed using tear sac massage and lid cleansing (neonatal lacrimal duct obstruction).
- **Red Flag**. Soft contact lens related infective conjunctivitis with corneal involvement. Make a habit of routinely asking whether soft contact lenses are worn. If available, use topical fluorescein to identify corneal staining (epithelial defect). Refer urgently if suspected corneal ulcer, and not give antibiotics in the interim as this may interfere with a subsequent corneal culture. Aim to have a low threshold for referral.

Differential

- Blepharoconjunctivitis and dry eye. Occurs when the acute episode is superimposed on longstanding intermittent lid redness, crusts, grittiness and watering. Best treated with lid hygiene (local heat, oil gland massage and lid

scrubs) and tear supplements until natural tear quality improves. Topical antibiotics do not have a place in treatment here.

- Atopic or allergic conjunctivitis. Recurrent seasonal or environment related red eye often with other history of atopy. Topical mast cell inhibitors and lid hygiene to reduce allergen load
- Toxic – drop allergy, preservative-related. Periorbital skin involvement, and often inferior medial conjunctiva worse affected in keeping with topical exposure
- Physiological - over use of drops that whiten conjunctiva for cosmetic reasons, leads to permanently dilated vessels and red eyes
- Contact lens related. Often signs in superior conjunctiva especially under upper lid. Stop contact lens use, avoid all preserved topical medication. If available, use fluorescein drops to check for corneal stain. Otherwise lid hygiene, non-preserved lubricant drops or mast cell stabiliser
- Intraocular causes of red eye usually affect vision and cause significant pain – eg acute glaucoma, scleritis, keratitis, uveitis (photophobia) and if severe signs refer for ophthalmic diagnosis and management
- Consider systemic associations e.g. thyroid eye disease

How to manage acute infective conjunctivitis in adult

- There is considerable discussion in the literature around prescribing topical antibiotics for a self-limiting condition. Patient education is probably the most important factor here.
- Delayed antibiotic prescribing for 3 days is the recommended strategy for conjunctivitis in primary care. This strategy is associated with reduced antibiotic use, with no evidence of medicalisation³
This trial showed that when GPs did not prescribe antibiotics at all for acute infective conjunctivitis, this still resulted in significant antibiotic use (30%) as patients would later re-present to the GP or buy over the counter topical antibiotics.
- A patient information leaflet may reduce time in consultation advising that symptoms remain the same but go on longer by only 1 day if no drops used.
- Infective conjunctivitis is best treated by bathing eyes with cool clean water, using lubricating drops, using moist cotton wool to clean the lids, and abstain from contact lens use.

How to manage acute infective conjunctivitis in children

- Even less evidence that drops effective at reducing symptom severity and duration of symptoms⁵

- Although official advice is that isolated cases (as opposed to outbreaks) of red eye do not need to stay away from school⁹ most nursery and primary schools still run an exclusion policy¹¹ advising same day attendance to GP and medication. This conflict needs to be resolved.

Key messages for commissioners

- A program of GP and patient education around eye health could be funded, eg. The use of effective lid hygiene practice, eye drop self-administration, and so on
- Effective and regular communication between GPs and local eye care professionals in high street, community clinics and hospital primary care services to develop protocols
- Involvement of local nurseries and primary schools in protocol implementation
- Topical antibiotic prescribing guidelines and feedback to individual practices

e-Learning for Health: <http://www.e-lfh.org.uk/projects/egp/index.html>

Useful Resources

- Public Health England Documents: Factsheet on Conjunctivitis. Conjunctivitis Information Leaflet 2013 Both available at http://www.hpa.org.uk/infections/topics_az/schools/ (accessed 2 February 2014)

People involved in creating this resource:

Miss Gillian Vafidis, Consultant Ophthalmologist
Dr Waqaar Shah, RCGP Clinical Champion in Eye Health

March 2013

References

1. Rietveld RP, ter Riet G, Bindels PJ, Sloos JH, van Weert HC. Predicting bacterial cause in infectious conjunctivitis: cohort study on informativeness of combinations of signs and symptoms. *BMJ* 2004;329:206-10.
2. Sheikh A, Hurwitz B. Topical antibiotics for acute bacterial conjunctivitis: Cochrane systematic review and meta-analysis update. *Brit J Gen Pract* 2005;55:962-4.
3. Everitt HA, Little PS, Smith PWF. A randomised controlled trial of management strategies for acute infective conjunctivitis in general practice. *BMJ* 2006;333:321.
4. [Rose PW](#) [Management strategies for acute infective conjunctivitis in primary care: a systematic review.](#) *Expert Opin Pharmacother.* 2007 Aug;8(12):1903-21.
5. Rose PW, Harnden A, Brueggemann AB, Perera R, Sheikh A, Crook D, et al. Chloramphenicol treatment for acute infective conjunctivitis in children in primary care: a randomised double-blind placebo-controlled trial. *Lancet* 2005;366:37-43.
6. [Sheikh A](#), [Hurwitz B](#), [van Schayck CP](#), [McLean S](#), [Nurmatov U](#). [Antibiotics versus placebo for acute bacterial conjunctivitis.](#) *Cochrane Database Syst Rev.* 2012 Sep 12;9:CD001211. doi: 10.1002/14651858.CD001211.pub3.
7. McCormick A, Fleming D, Charlton J. *Morbidity statistics from general practice. Fourth national study 1991-2.* London: HMSO, 1995.
8. Dart JK. Eye disease at a community health centre. *Br Med J (Clin Res Ed)* 1986; 293: 1477–1480.
9. Health Protection Agency. Guidelines on the management of communicable diseases in schools and nurseries: Conjunctivitis. Available at www.hpa.org.uk/infections/topics_az/schools/guideline_info/conjunctivitis.htm
10. Rose PW, Ziebland S, Harnden A, Mayon-White R, and Mant D. Why do General Practitioners prescribe antibiotics for acute infective conjunctivitis in children? Qualitative interviews with GPs and a questionnaire survey of parents and teachers. *Family practice* 2006; 23: 226–232.